

# Progressive Spinal Degeneration

## Phase Three



### **Incidence:**

Most often 40 to 65 years

### **Onset:**

Phase two, a vertebral subluxation that remained uncorrected.

### **Histopathology and X-ray:**

Beginning of bone fusion evident  
Massive advanced spur and arthritic formation.  
Abnormal calcium production  
Soft tissue decay is very evident  
Vertebral deformity has reached major proportions  
Disc function is lost and discs are degenerated  
Poor nerve function; neurological transmission is seriously impaired  
Nerve failure has started  
Muscular atrophy and wasting of shoulder and arm muscles

### **Most common symptoms:**

Severe pain (sometimes none at all)  
Greatly reduced movement  
General over-all stiffness  
Atrophy and wasting of shoulder and arm muscles  
Weakness in grip strength  
Fatigue  
Spells and dizziness  
Visual and ear disturbances  
Face numbness and pain  
Severe headaches  
Lowered resistance  
Chronic sinus problems  
Uncoordinated  
Irritability  
Vertebral artery insufficiency, which can lead to stroke  
Arm and hand pain and numbness

### **Corrective time:**

Usually 30 to 42 months

### **% of Recovery:**

Dependant on severity and patient compliance

## **Gooing Chiropractic Clinic**

2787 Bristol Suite 102

Costa Mesa, CA 92626

(714) 556-9188